



De Rham Cohomology for Compact Kahler Manifolds

Deep Bhattacharjee

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

July 31, 2023

De Rham Cohomology for compact Kähler manifolds

Deep Bhattacharjee

Abstract:

De Rham Cohomology is shown for compact Kähler manifolds considering the Hodge theory, Kähler potential and ddbar lemma.

Keywords: Kähler manifolds, Hodge Theory, ddbar Lemma.

MSC (2020): primary – 14-XX, secondary – 14F40

Introduction:

The ddbar lemma is a powerful tool whose bidegree form is essential in incorporating the De Rham Cohomology that in turn expresses the compact Kahler manifold provided certain conditions are met. Here in this paper the exact and harmonic form is taken along with the De Rham Cohomology class to compute the Kahler potential and ddbar lemma whose consequence is the Hodge theory.

Methodology:

Let there be an external mapping of sections for the exterior derivative Δ for the mapping of the projections that in general turned out to be the Dolbeault operators that in essence be the,

$$\Delta: \Omega^r \rightarrow \Omega^{r+1}$$

Where for the De Rham Cohomology one can describe the 4 forms as $\epsilon, \epsilon_1, \epsilon_2, \epsilon_3$ for a equivalence class of closed forms $[\epsilon]$ having the representation,

$$\epsilon_1 = \epsilon_2 + \epsilon_3$$

Where for $\epsilon_1 \cong [\epsilon]$, one can get the harmonic form $\Delta\epsilon_3 = 0$ for the exact form ϵ_2 . Thus, if we take a constant sheaf on $\overline{\mathcal{R}}$ for a smooth manifold S where the De Rham Cohomology showed that for any map parameterized by J one can easily make the mapping,

$$J : H_{dR}^p(S) \rightarrow H^p(S, \overline{\mathcal{R}}) \ni \begin{cases} [\epsilon] \in H_{dR}^p(S) \\ \forall J_{\epsilon_1} \in Hom H_p(S, \overline{\mathcal{R}}) \cong H^p(S, \overline{\mathcal{R}}) \end{cases}$$

This proves two identities;

1. One can have a natural isomorphism for the sheaf cohomology of $\overline{\mathcal{R}}$ in a way as to show,

$$H_{dR}^*(S) \cong H^*(S, \overline{\mathcal{R}}) \\ \exists \overline{\mathcal{R}} \text{ represents a Abelian group.}$$

2. One gets the isomorphism between the singular cohomology and de Rham cohomology such that for ant set \mathcal{G} and a trivial parameter classification or class $[G]$ there exists the relation with ϵ_1 in the way,

$$T \approx [G] \rightarrow \sum_G \epsilon_1 \quad \forall \begin{cases} T \in [G] \\ T \in H_p(S) \end{cases}$$

Thus, one can take the exact form ϵ_2 and differentiating it with i, j, k for a grouping of $\langle |i|, |j| \rangle$ the resultant factor provides the relation to $\Delta: \Omega^r \rightarrow \Omega^{r+1}$ such that in the case of ϵ_2 one of the most important aspects of Hodge theory can be found giving the wedge form,

$$\Omega^{p,q} \ni \epsilon_2 \cong \sum_{p,q} f_{ij} \Delta z^i \wedge \Delta \bar{z}^j$$

\exists in $\langle |i|, |j| \rangle; |i| = p, |j| = q$ for $\Omega^{p,q}$

Thus, one can find $\Delta = \partial + \bar{\partial}$ in differentiating the exact form ϵ_2 such that,

1. $\partial \epsilon_2$
2. $\bar{\partial} \epsilon_2$
3. $\partial \bar{\partial} + \bar{\partial} \partial = 0$
 - where Poincaré Lemma holds for ∂ and $\bar{\partial}$
 - for ϵ_2 in $\Delta \epsilon_2$ complex differential it is $\partial \bar{\partial}$ lemma

Thus, for the $\partial \bar{\partial}$ lemma, one can satisfy compact manifolds as Kähler provided in the consequence of Hodge Theory, if one corresponds, the $\Delta \epsilon_2$ norm then, for the compact Kähler, a global form of this lemma holds.

Let L be a compact Riemannian manifold, then for the relation: $\epsilon_1 = \epsilon_2 + \epsilon_3$. When $\Delta \epsilon_3 = 0$ then there exists exactly one ϵ_2 - form for the De Rham Cohomology class in $H_{dR}^K(L)$. Then for the space of the harmonic (ϵ_3) k - forms L is isomorphic to $H^k(L, \bar{\mathcal{R}})$ taking the sheaf Cohomology for K^{th} - Betti numbers in each of such finite spaces. Thus in this case it can be assumed that the manifold $L \cong S$ for the Abelian group of \mathcal{R} .

Therefore, taking the complex manifold (Kähler) K having a exact form $\Omega^{p,q} \ni \epsilon_2$ the $\partial\bar{\partial}$ lemma takes the bidegree (1,1) form of $\Omega^{p,q} \forall p, q \geq 1$ for a relation with the De Rham Cohomology such that in the k – forms one can get the exact form of the Kähler $\Omega^k(K)$ whose class is zero is De Rham Cohomology for $H_{dR}^{p+q}(K, \mathbb{C})$ has also the $\partial\bar{\partial} = 0$. This bidegree form is essential for the Kähler potential for $[\epsilon]$ such that in the case of the relation $\epsilon_1 \cong [\epsilon]$, the potential is defined,

$$\epsilon_1 = i^{-2}\partial\bar{\partial}\rho$$

Where for the Kähler manifold (K, ϵ_1) for the potential of Kähler to be defined as ρ for $[\epsilon]$, there exists the neighborhood μ of $\bar{\rho}$ where $\bar{\rho}$ is a local Kähler potential for the exact differential of (K, \mathbb{C}) such that, in compact forms of Kähler one can get the ϵ_2 form of the potential in the local potential $\bar{\rho}$ for the form in $\mu \subset K$,

$$i\partial\bar{\partial}(K, \mathbb{C}) \equiv \Omega^{1,1}(K)$$

For the $\partial\bar{\partial}$ –manifolds, when it has been assumed the compact space to be L previously, where the compact Kähler denoted as K then its not difficult to say that,

$$L \approx K$$

Conclusion:

For the bidegree form of the ddbar lemma; the most important consequences can be seen as the Kahler potential for the Kahler manifolds more specifically that manifold is compact. While taking the harmonic form and sheaf theoretic method into consideration, the related identities can be easily proved where at the end the manifold has been established as Kahler that also in compact form for the De Rham Cohomology.

References:

- Bhattacharjee, D., Roy, S. S., & Behera, A. K. (2022, August 31). *Relating Enriquer surface with K3 and Kummer through involutions and double covers over finite automorphisms on Topological Euler–Poincaré characteristics over complex K3 with Kähler equivalence*. <https://doi.org/10.21203/rs.3.rs-2011341/v1>
- Bhattacharjee, D. (2023, May 2). *Eliminating Hyperbolic Curvature of 9-Dimensions in Quotient Space for Elliptic Generators*. <https://doi.org/10.21203/rs.3.rs-2879172/v1>
- Bhattacharjee, D. (2022, May 10). *Establishing equivalence among hypercomplex structures via Kodaira embedding theorem for non-singular quintic 3-fold having positively closed (1,1)-form Kähler potential $\bar{\partial}^{-1}\partial\bar{\partial}^*\rho$* . <https://doi.org/10.21203/rs.3.rs-1635957/v1>
- Bhattacharjee, D. (2022, August 16). *Generalization of Quartic and Quintic Calabi – Yau Manifolds Fibered by Polarized K3 Surfaces*. <https://doi.org/10.21203/rs.3.rs-1965255/v1>
- Bhattacharjee, D. (2022, April 11). *Rigorously Computed Enumerative Norms as Prescribed through Quantum Cohomological Connectivity over Gromov – Witten Invariants*. <https://doi.org/10.36227/techrxiv.19524214.v1>
- Kunz, E. (2012, November 6). *Introduction to Commutative Algebra and Algebraic Geometry*. Springer Science & Business Media.
- Preuß, G. (2011, June 27). *Foundations of Topology*. Springer Science & Business Media.
- Putinar, M., & Sullivant, S. (2008, December 10). *Emerging Applications of Algebraic Geometry*. Springer Science & Business Media.
- Hatcher, A. (2002, January 1). *Algebraic Topology*. Cambridge University Press.
- Spanier, E. H. (1989, January 1). *Algebraic Topology*. Springer Science & Business Media.
- Bhattacharjee, D., Roy, S. S., Sadhu, R., & Behera, A. K. (2023, July 10). KK Theory and K Theory for Type II Strings Formalism. *Asian Research Journal of Mathematics*, 19(9), 79–94. <https://doi.org/10.9734/arjom/2023/v19i9701>